


Letz Creek Drainage



General location

General characteristics

7th field huc id = 17100206010303

Parent watershed = UPPER SIUSLAW RIVER

Total acreage = 5113

Maximum elevation = 464 feet

Minimum elevation = 176 feet

Ecological Capital

14 percent of the catchment has potential to contribute low to the aquatic system

39 percent of the stream system has adequate shading

40 percent of the riparian area is in good condition

16 miles of stream have inherently good coho spawning and rearing habitat

65 acres of potential or existing wetlands are present within the catchment


Potential Threats

There are 98 points where roads cross over fish bearing streams

Riparian road density = 0.84 miles per square mile

Mid-slope road density = 0.55 miles per square mile

1 percent of the catchment is considered to have a high potential of land slide occurrence




Ownership Patterns

4 percent of the catchment is private non-industrial

52 percent of the catchment is private industrial

42 percent of the catchment is federally owned

less than 1 percent of the catchment falls on other public lands



Notes

The catchment is dominated by moderate or low gradient confined or moderately confined channel habitat types. A total of 6.77 miles of stream are considered anadromous fish bearing and no miles of stream have digitized habitat surveys. A total of 3.90 miles of spawning surveys have been conducted since 1990 reflecting relatively moderate numbers of coho spawners. No snorkel surveys have been conducted in this catchment.

Due to lack of large diameter trees in the riparian area or directly contributing to the aquatic system, large woody debris in the stream system is most likely in short supply.

Stream temperatures may be high due to the high percent of streams exposed to direct sunlight. Streamside shading is most likely limiting water quality for fish habitat.

This catchment has low potential for consideration for anchor habitat status