


Lower Greenleaf Creek Drainage



General location

General characteristics

7th field huc id = = 171 00206040302

Parent watershed = LAKE CREEK

Total acreage = 4206

Maximum elevation = 700 feet

Minimum elevation = 115 feet

Ecological Capital

18 percent of the catchment has potential to contribute lwd to the aquatic system

51 percent of the stream system has adequate shading

42 percent of the riparian area is in good condition

7 miles of stream have inherently good coho spawning and rearing habitat

21 acres of potential or existing wetlands are present within the catchment

Potential Threats

There are 33 points where roads cross over fish bearing streams

Riparian road density = 0.92 miles per square mile

Mid-slope road density = 1.00 miles per square mile

4 percent of the catchment is considered to have a high potential of land slide occurrence


Ownership Patterns

4 percent of the catchment is private non-industrial

5 percent of the catchment is private industrial

69 percent of the catchment is federally owned

20 percent of the catchment falls on other public lands



 *Anadromous fish bearing streams*

 *Potential problem culverts*

 *In-stream habitat and wetland restoration / revegetation projects*

Notes

The catchment is dominated by moderate or low gradient confined or moderately confined channel habitat types. A total of 3.98 miles of stream are considered anadromous fish bearing and 3.98 miles of stream have digitized habitat surveys. A total of 4.38 miles of spawning surveys have been conducted since 1990 reflecting relatively high numbers of coho spawners. No snorkel surveys have been conducted in this catchment.

There is most likely sufficient lwd production and input into the aquatic system. The location of lwd sources should be identified and efforts should be made to maintain production.

Stream temperatures may be high due to the high percent of streams exposed to direct sunlight. Streamside shading is most likely limiting water quality for fish habitat.

Because of high fish productivity and high amounts of ecological capital this catchment has high potential to be an anchor habitat and should be evaluated in greater detail